

Global School-based Student Health Survey

Botswana 2005 Fact Sheet



The **2005 Botswana GSHS** measured alcohol and other drug use; sexual behaviors that contribute to HIV infection, other STI, and unintended pregnancy; unintentional injuries and violence; hygiene; dietary behaviors and overweight; physical activity; tobacco use; mental health; and protective factors.

The 2005 Botswana GSHS was a school-based survey of students in forms 1, 2, and 3. A two-stage cluster sample design was used to produce data representative of all students in forms 1, 2, and 3 in Botswana. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the student response rate was 95%, and the overall response rate was 95%. A total of 2,197 students participated in the Botswana GSHS. Students self-reported their responses to each question on a computer scannable answer sheet. For comparison purposes, only students aged 13-15 years are included in the analyses for this fact sheet.

| Results for students aged 13-15 years | Total | Boys | Girls |
|--|------------|------------|------------|
| Alcohol and Other Drug Use | | | |
| Percentage of students who had at least one drink containing alcohol on one or more days during the past 30 days | 20.6 ± 2.6 | 22.8 ± 3.6 | 18.7 ± 2.9 |
| Percentage of students who drank so much alcohol that they were really drunk one or more times during their life | 20.9 ± 2.1 | 24.9 ± 3.4 | 17.4 ± 2.2 |
| Percentage of students who had a hang-over, felt sick, got into trouble with their family or friends, missed school, or got into fights, as a result of drinking alcohol one or more times during their life | 16.6 ± 2.2 | 19.2 ± 3.9 | 14.5 ± 2.2 |
| Percentage of students who used drugs one or more times during their life | 7.5 ± 1.6 | 13.0 ± 2.6 | 2.8 ± 1.4 |
| Sexual Behaviors That Contribute to HIV Infection, Other STI, and Unintended Pregnancy | | | |
| Percentage of students who have ever had sexual intercourse | 17.4 ± 3.0 | 26.9 ± 5.1 | 10.3 ± 2.5 |
| Percentage of students who have had sexual intercourse with two or more people during their life | 10.0 ± 2.4 | 15.5 ± 3.6 | 5.8 ± 2.5 |
| Unintentional Injuries and Violence | | | |
| Percentage of students who were physically attacked one or more times during the past 12 months | 55.7 ± 4.3 | 56.9 ± 4.6 | 54.7 ± 5.2 |
| Percentage of students who were in a physical fight one or more times during the past 12 months | 47.7 ± 3.4 | 54.4 ± 5.0 | 41.7 ± 3.6 |
| Percentage of students who were seriously injured one or more times during the past 12 months | 65.8 ± 3.3 | 68.6 ± 5.0 | 63.3 ± 3.3 |
| Percentage of students who were bullied on one or more days during the past 30 days | 52.1 ± 3.7 | 52.6 ± 4.9 | 51.8 ± 3.8 |

Global School-based Student Health Survey

Botswana 2005 Fact Sheet



| Results for students aged 13-15 years | Total | Boys | Girls |
|---|------------|------------|------------|
| Hygiene | | | |
| Percentage of students who cleaned or brushed their teeth less than 1 time per day during the past 30 days | 5.1 ± 1.5 | 7.3 ± 1.8 | 3.0 ± 1.8 |
| Percentage of students who never or rarely washed their hands before eating during the past 30 days | 4.5 ± 1.4 | 7.0 ± 2.3 | 2.4 ± 1.4 |
| Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days | 5.2 ± 1.0 | 6.9 ± 1.4 | 3.8 ± 1.5 |
| Dietary Behaviors and Overweight | | | |
| Percentage of students who went hungry most of the time or always during the past 30 days because there was not enough food in their home | 14.2 ± 2.3 | 15.9 ± 3.3 | 12.8 ± 3.2 |
| Physical Activity | | | |
| Percentage of students who were physically active for a total of at least 60 minutes per day on all 7 days during the past 7 days | 11.7 ± 1.4 | 13.0 ± 3.0 | 10.4 ± 2.2 |
| Percentage of students who spent three or more hours per day during a typical or usual day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities | 34.5 ± 2.8 | 35.3 ± 2.7 | 33.7 ± 4.1 |
| Tobacco Use | | | |
| Percentage of students who smoked cigarettes on one or more days during the past 30 days | 7.0 ± 1.4 | 10.7 ± 2.6 | 3.8 ± 1.5 |
| Percentage of students who used any other form of tobacco on one or more days during the past 30 days | 11.6 ± 1.4 | 14.0 ± 1.5 | 9.4 ± 2.3 |
| Percentage of students who reported people smoking in their presence on one or more days during the past 7 days | 45.0 ± 3.3 | 47.5 ± 4.2 | 42.7 ± 4.4 |
| Mental Health | | | |
| Percentage of students who felt lonely most of the time or always during the past 12 months | 16.5 ± 2.6 | 15.6 ± 3.3 | 17.1 ± 3.6 |
| Percentage of students who seriously considered attempting suicide during the past 12 months | 22.4 ± 2.8 | 20.5 ± 4.2 | 23.8 ± 3.9 |
| Percentage of students who have no close friends | 14.0 ± 2.2 | 13.5 ± 2.3 | 14.6 ± 3.2 |

Global School-based Student Health Survey



Botswana 2005 Fact Sheet

| Results for students aged 13-15 years | Total | Boys | Girls |
|---|------------|------------|------------|
| Protective Factors | | | |
| Percentage of students who missed classes or school without permission on one or more days during the past 30 days | 23.3 ± 3.4 | 28.7 ± 4.2 | 18.9 ± 4.1 |
| Percentage of students who reported that most of the students in their school were never or rarely kind and helpful during the past 30 days | 27.7 ± 3.0 | 32.2 ± 4.9 | 23.9 ± 3.1 |
| Percentage of students whose parents or guardians never or rarely really knew what they were doing with their free time during the past 30 days | 31.6 ± 2.8 | 34.3 ± 3.4 | 29.3 ± 3.9 |

For additional information, please contact:

Mrs. Rachel Mosinyi, Ag. Principal Health Officer (HE)
Ministry of Health, Gaborone, Botswana, r_mosinyi@yahoo.co.uk